

Disinfectant and cleaner for use on non-invasive devices, equipment and surfaces in medical, surgical and dental health care facilities such as stethoscopes, stainless steel or plastic trays and alligator clips.

GamaCide effectively kills the following microorganisms at room temperature (20°C) with a 1 minute contact time: *Mycobacterium terrae* ATCC 15755, *Staphylococcus aureus* ATCC 6538, *Salmonella choleraesuis* ATCC 10798, *Pseudomonas aeruginosa* ATCC 15442, *Escherichia coli* NCTC 8196, *Enterococcus hirae* ATCC 10541, Poliovirus 1 strain Chat ATCC VR-1562 and Influenza A viruses. Effectively inactivates SARS-Cov-2 (COVID 19), Herpes Simplex 1 virus, Human immunodeficiency virus type 1, Human hepatitis B virus in 2 minutes. A 3-minute contact time is required to kill *Trichophyton mentagrophytes* ATCC9533.

DIRECTIONS: Appropriate glove protection is recommended prior to use of product. Use where control of the hazards of cross-contamination is of concern. Prior to surface disinfection, clean soiled surfaces by saturating surface with GamaCide[®] solution allowing surfaces to remain wet for 30 seconds, then wipe clean using a fresh paper or cloth towel. Repeat the cleaning process until surfaces are visibly clean. Make sure surfaces are free of all gross filth and heavy soil. Discard paper towels or launder cloth towels before reusing. **DO NOT SWALLOW.**

GENERAL DISINFECTION: Apply GamaCide[®] solution to surface. Allow surface to remain wet for 3 minutes at room temperature (20°C) for broad spectrum disinfection then rinse or wipe dry using a clean paper or cloth towel. Discard paper towels, or launder cloth towel before reusing. Surgical instruments must be sterilized according to manufacturer's instructions after cleaning with GamaCide[®]. Do not mix with detergents or other chemicals. Use undiluted. Do not reuse the same solution after application on medical device(s).

PERSONAL PROTECTION: Wear appropriate barrier protection such as gloves, gowns, masks or eye coverings. Cleaning Procedure: Blood and other body fluids must be thoroughly cleaned from surfaces/objects before disinfection. Wipe the pre-cleaned area to be disinfected. Allow surface to remain wet for 3 minutes for broad spectrum disinfection.

ULTRASONIC CLEANING SOLUTION/ MANUAL CLEANER: Thoroughly pre-rinse objects under running water to remove visibly gross debris. Immerse instruments in solution and activate ultrasonic unit for a minimum of 1 minute. For manual cleaning, immerse instruments in solution and scrub using a stiff bristle brush until visibly clean (submerge objects while scrubbing). Rinse thoroughly. Change solution as it becomes diluted or visibly soiled.

CONTRAINDICATIONS: Not for use in hospital nurseries or on children's toys. Professional use only. Do not use to disinfect invasive devices. Invasive devices must be disinfected or sterilized prior to use.

GERMIPHENE GamaCide³_{TM}

Disinfectant for Non-Invasive Devices, Equipment, and Surfaces
Bactericidal • Fungicidal • Virucidal • Tuberculocidal

Medical/Surgical/Dental Disinfectant Cleaner
For Professional Use Only. Ready to Use.

ACTIVE INGREDIENTS:

Benzethonium Chloride.....0.35% w/w
Isopropyl Alcohol.....20.00% w/w

4L

MD REF EU-GAMA-XXL

UDI



(01) 0 0850046 43635 2
(10) LOT123
(17) 260309

LOT ???????



YYYY-MM-DD

INFECTIOUS MATERIALS DISPOSAL: Cleaning materials used that may contain blood/body fluids should be autoclaved and/or disposed of in accordance with local regulations for infectious materials disposal.

STORAGE AND DISPOSAL: Do not contaminate food or feed by storage or disposal. Storage: Store in cool, well ventilated area. Product Disposal: Unused contents to be disposed of in accordance with local/regional/national regulations. Container Disposal: Triple rinse empty container, then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill.



FLAMMABLE: Do not use in presence of open flame or spark.
CAUTION: KEEP OUT OF REACH OF CHILDREN. Warning! Causes eye and skin irritation. Avoid contact with eyes, skin, or clothing. Wear protective gloves, clothing, goggles, face protection. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse. For prolonged, frequent or continued use of more than 15 minutes, wear protective gloves, eye wear and face protection. Avoid contact with skin and eyes. Keep from open flame or spark. Flammable liquid and vapor. Do not use in presence of open flame or spark. Keep container tightly closed.

FIRST AID: For more information, refer to SDS. **IF ON SKIN:** Take off contaminated clothing. Rinse skin immediately with water. **IF IN EYES:** Rinse eyes with water for several minutes. Remove contacts, if worn, and continue rinsing until advised to stop by the Poisons Information Centre or a doctor, or for at least 15 minutes. If skin irritation occurs or eye irritation persists, get medical attention. Have the product container/label with you when calling a Poison Control Centre or doctor, or going for treatment.



Young Microbrush Ireland Ltd.
Clogherane, Dungarvan
Co. Waterford, X35 VEOz, Ireland



EU-GM3-LB-4L R1



CEpartner4U
Esdoornlaan 13, 3951 DB Maam, Netherlands



Manufactured by: **Germiphene Corporation**
1379 Colborne St. E. | Brantford ON | Canada N3T 5M1
Toll Free Phone 1.800.265.9931



For a list of additional translations, please go to:
<https://germiphene.com/product-resources/>



MADE IN CANADA

4 in

10 in