





Disinfectant for Non-Invasive Devices, Equipment, and Surfaces Bactericidal • Fungicidal • Virucidal • Tuberculocidal Medical/Surgical/Dental Disinfectant Cleaner For Professional Use Only. Ready to Use.

## ACTIVE INGREDIENTS:

Benzethonium Chloride. ...0.35% w/w Isopropyl Alcohol. 

Disinfectant and cleaner for use on non-invasive devices Disinfectant and cleaner for use on non-invasive devices, equipment and surfaces in medical, surgical and dental health care facilities such as stethoscopes, stainless steel or plastic trays and alligator clips.

GamaCide effectively kills the following microorganisms at room GamaCide effectively kills the following microorganisms at roo temperature (20 C) with a 1 minute contact time: Mycobacteriu terrae ATCC 15755, Staphylococcus aureus ATCC 6538, Salmonelle cholereasus ATCC 15798, Pseudomonas aeruginosa ATCC 15424, Escherichia coli NCTC 8396, Enterococcus hirae ATCC 15941, Poliovirus 1 strain Chat ATCC VR-1562 and Influenza A viruses. Foroving Fiscality Classifier Cover (COVID) (c), Herpes Simplex 1 Virus, Human immunodeficiency virus type 1, Human hepatitis B virus in 2 minutes. A 3-minute contact time is required to kill Trichophyton mentagrophytes ATCC9533.

DIRECTIONS: Appropriate glove protection is recommended prior to use of product. Use where control of the hazards of cross-contamiation is of concern. Remove large cover of canister. Remove top of refill pouch and place pouch in canister. Locate first when in center of the roll and thread through eyelet of cover. Replace cover. Pull each wipe out at an angle from the cap. Close cap after each use. DO NOT SWALLOW.

**GENERAL DISINFECTION:** Thoroughly clean the surface of heavy and organic soil prior to disinfection. Wipe the pre-cleaned area to be disinfected. Allow surface to remain wet for 3 minutes for broad spectrum disinfection. Surgical instruments must be sterilized according to manufacturer's instructions after cleaning with GamaCide<sup>3</sup> wipes.

PERSONAL PROTECTION: Wear appropriate barrier protection such as gloves, gowns, masks or eye coverings. Cleaning Procedure: Blood and other body fluids must be thoroughly cleaned from surfaces/ objects before disinfection.



Wipe the pre-cleaned area to be disinfected. Allow surface to remain wet for 3 minutes for broad spectrum disinfection

MD

IREF

EU-GAMA-SW-R

INFECTIOUS MATERIALS DISPOSAL: Cleaning materials used https://www.contain.blood/body/fluids.should be autoclaved and/or disposed of in accordance with local regulations for infectious materials disposal. Storage and Disposal: Do not contaminate materials disposal. Storage and Disposal. Do not contaminate food or feed by storage or disposal. Storage: Store in cool, well ventilated area. Product Disposal. Storage: Store in cool, well ventilated area. Product Disposal. Dispose of vipe in trash after use. Unused contents to be disposed of in accordance with local/regional/national regulations. Container Disposal: Triple rinse empty container, then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill.

CONTRAINDICATIONS: Not for use in hospital nurseries or on children's toys. Professional use only. Do not use to disinfect invasive devices. Invasive devices must be disinfected or sterilized prior to use.

## FLAMMABLE: Do not use in presence of KEEP OUT OF REACH OF CHILDREN. FLAMMABLE: Do not use in presence of open flame or spark

Warning! Causes eye and skin irritation. Avoid contact with eyes, skin, or clothing. Wear protective gloves, clothing, goggles, face protection. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

FIRST AID: For more information, refer to SDS. IF ON SKIN: Take off contaminated clothing. Rinse skin immediately with water. IF IN EYES: Rinse eyes with water for several minutes. Remove IN ETEX: kinse eyes with water for several minutes, kensore contacts, if worn, and continue rinsing until advised to stop by the Poisons Information Centre or a doctor, or for at least 15 minutes. If skin irritation occurs or eye irritation persists, get medical attention. Have the product container/label with you when calling a Poison Control Centre or doctor, or going for treatment.





CE Young Microbrush Ireland Ltd. 0050 Clogherane, Dungarvan Co. Waterford, X35 VE02, Ireland



EU-GM3-LB-SWR R1